

AGROSTEMIN[®]



Dr. Danilo Gajić



**EFEKAT FOLIJARNE PRIMENE AGROSTEMIN[®]-a
NA ZASAD DINJE**
– fenološka opažanja –
(*Cucumis melo*)



Ogledi su izvedeni u gradovima Mossoró i Baraúnas – RN Brazil.

AGROSTEMIN[®] je primenjen putem sistema za zalivanje (eng: "fertigation") između 5. i 7. dana nakon rasađivanja, u količini 30g po hektaru.

FENOLOŠKA OPAŽANJA

(fotografije)

PRVI OGLED

Sorta: GALIA

Hibrid: ESTORIL

Tretirano AGROSTEMIN® – **om:** *5 dana nakon rastađivanja*

STAROST biljaka: 21 dana od rastađivanja i 16 dana nakon tretiranja.



KONTROLA

TRETIRANO



bez
AGROSTEMIN®-a

sa
AGROSTEMIN®-om

DRUGI OGLED

Sorta: AMARELO

Hibrid: GOLD EX

Tretirano **AGROSTEMIN[®]**–om: *7 dana nakon rastađivanja*

STAROST biljaka: *23 dana od rastađivanja i 16 dana nakon tretiranja*

bez AGROSTEMIN®-a

sa AGROSTEMIN®-om



KONTROLA

TRETIRANO



KONTROLA

TRETIRANO

TREĆI OGLED

Sorta: GALIA

Varijetet: DENNY

Tretirano AGROSTEMIN[®] – om: *5 dana nakon rastađivanja*

STAROST biljaka: *22 dana od rastađivanja i 15 dana nakon tretiranja*





NETRETIRANO

TRETIRANO

ČETVRTI OGLED

Sorta: GALIA

Varijetet: 8530

Tretirano AGROSTEMIN[®]-om: *5 dana nakon rastađivanja*


STAROST biljaka: *20 dana od rastađivanja i 15 dana nakon tretiranja*



bez
AGROSTEMIN®-a

sa
AGROSTEMIN®-om





AGROSTEMIN[®]



Dr. Danilo Gajić

www.agrostemin.com